

CPD SAJCH May 2008

- 1. Which of the following is FALSE?
 - a) hepatitis B may be transmitted though blood products
 - b) hepatitis A is usually transmitted through the faeco-oral route
 - c) hepatitis A is rare but life threatening
 - d) hepatitis A is usually more common in children than adults
 - e) in fulminant hepatitis treatment with vitamin K can be life saving.
- 2. Which of the following is FALSE? Viral hepatitis may present with
 - a) jaundice
 - b) prolonged bleeding
 - c) encaphalopathy
 - d) pale urine
 - e) right upper quandrant pain.
- 3. True (A) or false (B) fill in either block A or block B: Viral hepatitis is not a notifiable disease.
- 4. True (A) or false (B) fill in either block A or block B: Viral hepatitis is usually symptomatic.
- 5. True (A) or false (B) fill in either block A or block B: Immunisation is likely to reduce the incidence of hepatitis A.
- 6. Which of the following is FALSE? Regarding noise levels
 - a) there are established standards of noise levels in work environments
 - b) some neonatal ICUs have levels well above those acceptable for adults
 - c) humans (ICU staff) may contribute more to noise levels than equipment
 - d) ICU noise is unlikely to cause sensorineural hearing impairment
 - e) the neonatal environment may influence later learning.
- 7. Which of the following is FALSE? Sialorrhoea
 - a) is the unintentional overflow of saliva from the oral cavity
 - b) is common in babies
 - c) is common in patients with neurological impairment
 - d) always requires surgical management
 - e) can lead to skin inflammation or infection.
- 8. True (A) or false (B) fill in either block A or block B: The main secretory control of the salivary glands is the sympathetic nervous system.
- 9. Which of the following is FALSE? Sialorrhoea
 - a) is a complex developmental brain disorder
 - b) ideally should be managed by a multidisciplinary team
 - c) can be worsened by nasal airway obstruction and maxillofacial abnormalities
 - d) can respond to botulinum toxin which reduces saliva production
 - e) can be treated using various surgical procedures.

SAJCH MAY 2008 VOL.2 NO.2

- 10. True (A) or false (B) fill in either block A or block B: Acute respiratory infections and gastroenteritis are the most common recorded causes of death in South African children.
- 11. True (A) or false (B) fill in either block A or block B: Almost half of recent deaths recorded in South African children had underlying advanced HIV disease.
- 12. True (A) or false (B) fill in either block A or block B: Child PIP is a retrospective mortality review process that is intended to inform and improve the quality of care that children receive.
- 13. With regards to Castleman's disease, which of the following is FALSE?
 - a) it is a lymphoproliferative disorder
 - b) the localised variant is often associated with constitutional symptoms
 - c) it usually presents as enlarged lymph nodes
 - d) it is usually associated with human herpesvirus 8 infection
 - e) complete excision is curative in the localised form.
- 14. Which of the following is NOT one of the aims of child PIP?
 - a) identification of all inpatient deaths
 - b) assigning a cause to each death
 - c) determining only the HIV status of each child who dies
 - d) determining factors that could have changed the outcome for each child who dies.
- 15. True (A) or false (B) fill in either block A or block B: Child PIP data are unable to describe information on the quality of health care provided.
- 16. True (A) or false (B) fill in either block A or block B: Child PIP data describe all paediatric deaths in South Africa.
- 17. True (A) or false (B) fill in either block A or block B: Child PIP is the initiative of the South African National Department of Health.
- 18. True (A) or false (B) fill in either block A or block B: There is adequate evidence for the effective treatment of multicentric angiofollicular lymph node hyperplasia associated with HIV infection with etoposide and vinblastine.
- 19. True (A) or false (B) fill in either block A or block B: Human herpesvirus 8-associated tumours tend to be very vascular in nature.
- 20. True (A) or false (B) fill in either block A or block B: Imaging soft-tissue masses can identify vascular lesions prior to surgery.

CPD questionnaires must be completed online via www.cpdjournals.org.za.

After submission you can check the answers and print your certificate.

Questions may be answered up to 6 months after publication of each issue.